

# High-field MRI reveals an acute impact on brain function in survivors of the magnitude 8.0 earthquake in China

Su Lui<sup>a</sup>, Xiaoqi Huang<sup>a,b</sup>, Long Chen<sup>a</sup>, Hehan Tang<sup>a</sup>, Tijiang Zhang<sup>a</sup>, Xiuli Li<sup>a</sup>, Dongming Li<sup>a</sup>, Weihong Kuang<sup>b</sup>, Raymond C. Chan<sup>c</sup>, Andrea Mechelli<sup>d</sup>, John A. Sweeney<sup>e</sup>, and Qiyong Gong<sup>a,f,1</sup>

<sup>a</sup>Huaxi MR Research Center, Department of Radiology, and <sup>b</sup>Department of Psychiatry, State Key Laboratory of Biotherapy, West China Hospital of Sichuan University, Chengdu 610041, China; <sup>c</sup>Institute of Psychology, Neuropsychology, and Applied Cognitive Neuroscience Laboratory, and Institute of Psychology, Key Laboratory of Mental Health, Chinese Academy of Sciences, Beijing 10001, China; <sup>d</sup>Section of Neuroimaging, Division of Psychological Medicine and Psychiatry, Institute of Psychiatry, King's College London, London SE5 8AF, United Kingdom; <sup>e</sup>Center for Cognitive Medicine, University of Illinois, Chicago, IL 60612; and <sup>f</sup>Division of Medical Imaging, Faculty of Medicine, University of Liverpool, Liverpool L69 3BX, United Kingdom

Edited by Marcus E. Raichle, Washington University School of Medicine, St. Louis, MO, and approved July 27, 2009 (received for review December 17, 2008)

Besides the enormous medical and economic consequences, national disasters, such as the Wenchuan 8.0 earthquake, also pose a risk to the mental health of survivors. In this context, a better understanding is needed of how functional brain systems adapt to severe emotional stress. Previous animal studies have demonstrated the importance of limbic, paralimbic, striatal, and prefrontal structures in stress and fear responses. Human studies, which have focused primarily on patients with clinically established posttraumatic stress disorders, have reported abnormalities in similar brain structures. At present, little is known about potential alterations of brain function in trauma survivors shortly after traumatic events. Here, we show alteration of brain function in a cohort of healthy survivors within 25 days after the Wenchuan earthquake by a recently discovered method known as “resting-state” functional MRI. The current investigation demonstrates that regional activity in frontolimbic and striatal areas increased significantly and connectivity among limbic and striatal networks was attenuated in our participants who had recently experienced severe emotional trauma. Trauma victims also had a reduced temporal synchronization within the “default mode” of resting-state brain function, which has been characterized in humans and other species. Taken together, our findings provide evidence that significant alterations in brain function, similar in many ways to those observed in posttraumatic stress disorders, can be seen shortly after major traumatic experiences, highlighting the need for early evaluation and intervention for trauma survivors.

neuroimaging | stress | anxiety | depression | posttraumatic stress disorder

In the afternoon on May 12, 2008, the epicenter of a devastating earthquake occurred in Wenchuan, in the Sichuan Province of China. It measured 8.0 on the Richter scale. The most severely affected geographical regions were Yingxiu, Wenchuan, Dujiangyan, and Shifang, where 45 million people were directly affected. Among them, 69,146 people were confirmed dead, 374,131 were seriously injured, and 17,516 are missing. A significant proportion of the survivors ( $\approx 20\%$ ) (1) are likely to develop stress-related disorders, such as acute stress disorder (ASD) and posttraumatic stress disorder (PTSD). Given the serious and persistent impact of these highly prevalent psychiatric disorders, it is vital to develop a better understanding of the alterations of cerebral function evident in the early stages of adaptation to trauma. Such knowledge may lead to a better understanding of posttraumatic responses and the development of more effective early interventions.

Studies of animal models of acute and chronic stress have provided evidence of physiological and morphological changes in several brain regions, including the amygdala, hippocampus, and prefrontal cortex (2, 3). Human studies (4–12) have focused primarily on patients who already have an established psychiatric

disorder, such as ASD and PTSD. Despite some inconsistencies (6, 7, 9–11), findings from clinical studies also implicate limbic, paralimbic, striatal, and prefrontal structures in the pathophysiology of stress-related psychiatric disorders. However, because the majority of these studies were performed with victims of trauma that took place years or even decades earlier, it is not clear whether this represents a slowly evolving pattern of brain alteration or one that emerges shortly after traumatic experiences. No study has yet investigated alterations in cerebral function in survivors soon after a massive widespread disaster, such as an earthquake.

We applied a recently discovered method known as “resting-state” functional MRI (fMRI) to assess brain function in a cohort of healthy survivors after the Wenchuan earthquake. “Resting-state” fMRI not only avoids performance confounds (13), but in the context of clinical studies to assess resting-state brain physiology it is easier to implement than positron emission tomography (PET)/single photon emission computed tomography (SPECT) because of its lower cost, greater availability, and noninvasiveness (14). Low-frequency (0.01–0.08 Hz) fluctuations of the blood-oxygenation-level-dependent (BOLD) signal in resting state fMRI data are thought to reflect spontaneous neural activity in nonhumans (15, 16) and humans (17, 18). In addition to regional amplitude of low-frequency (0.01–0.08 Hz) fluctuations (ALFF), resting-state fMRI provides data about functional connectivity. Although ALFF provides information about synchronous regional cerebral activity, analysis of cross-correlations between spatially remote regions allows the integrity of distributed brain networks to be examined.

In the present investigation, we applied resting-state fMRI to examine both regional cerebral function and functional brain connectivity in physically healthy trauma survivors shortly (13–25 days) after a massive psychological trauma. We hypothesized that survivors would show: (i) altered brain function shortly after the earthquake, especially in brain regions known to be important for emotion processing; and (ii) that alterations in brain function would be associated with survivors’ reports of emotional distress.

## Results

We acquired whole-brain resting-state fMRI in 44 healthy survivors and 32 healthy controls during resting state on a 3-T

Author contributions: S.L., X.H., and Q.G. designed research; S.L., X.H., L.C., H.T., T.Z., X.L., D.L., W.K., and Q.G. performed research; S.L. and L.C. analyzed data; and S.L., R.C.C., A.M., J.A.S., and Q.G. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. E-mail: qiyonggong@hmrc.org.cn.


Characteristics	Survivors $\pm$ SD ( <i>n</i> = 44)	Controls $\pm$ SD ( <i>n</i> = 32)	<i>P</i>
Female to male, no.	17:27	12:20	0.90
Mean age, y	37 $\pm$ 10.6	34.6 $\pm$ 11.0	0.34
Years of education	8.6 $\pm$ 4.1	9.3 $\pm$ 4.2	0.42
Days after earthquake	21.2 $\pm$ 3.3	—	—
SAS scores	48.4 $\pm$ 11.4	—	—
SDS scores	46.8 $\pm$ 10.8	—	—

To explore the alteration of regional activity in the survivor group, we examined the ALFF (0.01–0.08 Hz) of BOLD signals. ALFF maps in the survivor and control groups were compared on a voxelwise basis by using a two sample *t* tests in Statistical Parametric Mapping software (SPM2; Wellcome Department of Imaging Neuroscience; [www.fil.ion.ucl.ac.uk](http://www.fil.ion.ucl.ac.uk)). All of the analyses of whole-brain image data kept experimentwise type I error rates protected at  $P < 0.05$ , with a familywise error (FWE) correction procedure that considers the intercorrelation of the data structure. The whole-brain analysis indicated that, compared with controls, survivors showed significantly increased ALFF in the left prefrontal cortex and the left precentral gyrus, extending medially to the left presupplementary motor area (pre-SMA;  $P < 0.05$ ; Fig. 1 and Table 2). In addition, region of interest (ROI) analyses revealed significantly increased ALFF in bilateral insula and caudate and the left putamen in the survivor

To characterize the alteration of neural networks involving brain areas associated with emotion processing and memory (5) and the core areas of the default mode (19), functional connectivity analysis was performed by using a seed voxel correlation approach (see *Materials and Methods*). Connectivity maps were compared across survivors and controls by using a two-sample *t* test in SPM2. Analysis of the combined sample of survivors and controls showed a significant positive association among seed regions, mainly involving the bilateral amygdala, hippocampus, caudate, putamen, insula, parahippocampus, prefrontal cortex, cingulate cortex, and cerebellum ( $P < 0.05$ ). However, comparison of the two groups revealed that survivors had decreased functional connectivity compared with controls within a distributed network that included the bilateral amygdala, hippocampus, caudate, putamen, insula, anterior cingulate cortex, and cerebellum ( $P < 0.05$ ; Fig. 2). Functional connectivity involving default-mode areas was also decreased significantly in the survivor relative to the control group. No increased functional connectivity was found in the survivors relative to the controls in any seed region. These results suggest that the survivors expressed decreased temporal synchronization of resting neurophysiological activity among limbic and striatal areas, as well as within components supporting default-mode function.

We also examined (i) the relationship between regional hyperactivation and decreased connectivity and (ii) the relationship between connectivity and the level of emotional distress as measured by SAS and SDS scores. These analyses revealed no significant correlations between decreased connectivity and increased regional ALFF values, or between connectivity and level of emotional distress ( $P > 0.05$ ).

This study characterizes the alteration of distributed brain function at rest in a cohort of survivors of the Wenchuan earthquake in China within 25 days (range 13–25 days; mean  $\pm$  SD: 21  $\pm$  3) after the initial seismic event. Compared with controls, survivors showed hyperactivity of prefrontal-limbic and striatal systems, including left prefrontal cortex, pre-SMA, bilateral insula, bilateral caudate, and left putamen, as well as attenuated functional connectivity involving limbic-striatal areas and default-mode areas. The increased regional activity and reduced functional connectivity in frontolimbic and striatal regions occurred in areas known to be important for emotion processing (22). Atypical function in these areas has been implicated in previous studies of patients with chronic stress-



**Table 2. Voxel-based analysis of global ALFF in trauma survivors and healthy controls**

Location	Talairach			Voxel size, mm <sup>3</sup>	<i>P</i>
	<i>x</i>	<i>y</i>	<i>z</i>		
Left hemisphere: survivors > controls					
Precentral gyrus*	−56	12	5	2646	<0.001 <sup>†</sup>
Prefrontal cortex	−3	12	63	1890	0.0015 <sup>†</sup>
Caudate	−12	14	−3	243	0.026 <sup>‡</sup>
Insula	−39	14	−11	270	0.039 <sup>‡</sup>
Putamen	−15	14	−3	378	0.009 <sup>‡</sup>
Right hemisphere: survivors > controls					
Caudate	12	12	8	243	0.045 <sup>‡</sup>
Insula	42	9	2	351	0.013 <sup>‡</sup>

\*Including the pre-SMA.

<sup>†</sup>Type I error rate protected at  $P < 0.05$  with FWE correction.<sup>‡</sup>Type I error rate corrected by using SVC.

related disorders (5, 6, 8, 12, 23) and was related to the reported level of emotional distress in our sample. Thus, the present investigation expands understanding of stress-related changes of brain function in humans in two important ways. First, our study demonstrates that brain function in trauma survivors is altered within 25 days of a significant traumatic event. Second, our study provides evidence that alterations in brain function are not limited to focal changes in brain activity, but are also expressed at the level of altered functional integration within distributed limbic-striatal and default-mode networks. Furthermore, the decreased functional connectivity was not significantly related to regional ALFF alteration or level of emotional distress, suggesting that it may be related to different neurophysiological alterations and potentially may have a different functional impact.

Prefrontal-limbic and striatal systems and, more recently, the pre-SMA, have been recognized to be involved in affective processing (24) and decision making (25). Functional neuroimaging studies have provided direct evidence that prefrontal-limbic and striatal systems play a critical role in anxiety disorders, including the recollection of traumatic memories and the processing of fear and pain (6, 26), whereas striatum and pre-SMA are activated when making decisions under time pressure (25). Studies in both humans (27, 28) and animals (29) also indicate the roles of these systems in regulation of stress-induced activation of the hypothalamic–pituitary–adrenal axis. Overactivity within these systems has been reported with PTSD patients in several fMRI studies using a range of experimental tasks (5, 6, 12, 26), and also with PET/SPECT studies at resting state (5).


**Table 3. Relationship of ALFF values with anxiety (SAS) and depression (SDS) scores of earthquake trauma survivors**

Regions	SAS, $r$ ( $P$ )	SDS, $r$ ( $P$ )
Left		
Amygdala	0.28 (0.07)	0.24 (0.12)
Caudate	0.23 (0.15)	0.23 (0.14)
Hippocampus	0.24 (0.13)	0.19 (0.22)
Insula	0.25 (0.11)	0.15 (0.33)
Putamen	0.47 (0.002)*	0.42 (0.005)*
Prefrontal	0.03 (0.86)	−0.08 (0.62)
Precentral gyrus	0.11 (0.49)	0.16 (0.32)
Right		
Amygdala	0.40 (0.009)*	0.44 (0.004)*
Caudate	0.37 (0.013)*	0.38 (0.013)*
Hippocampus	0.39 (0.011)*	0.35 (0.024)*
Insula	0.10 (0.52)	0.08 (0.63)
Putamen	0.29 (0.059)	0.25 (0.12)

\* $P < 0.05$  with Bonferroni correction.

Our findings in the survivor group showing increased activity in the resting state in the same set of areas support our hypothesis that regional cerebral function can be affected in trauma survivors shortly after they experience severe traumatic events. This is an important finding because it indicates that changes observed in patients with chronic stress-related disorders do not only reflect slowly progressive alterations or secondary effects of chronic distress and disability. Rather, dysfunction in clinically relevant brain circuitry seems to be already manifested shortly after massive traumatic events. Consistent with this idea, studies based on animal models have revealed that cortical GABA neurons can be modulated by allopregnanolone soon (3–4 weeks) after subjecting animals to experimental fearful stimuli (30). The increased GABA tone could increase regionally synchronized neuronal activity (31) and activity in  $\gamma$ -band power (32), which has been found to be correlated with ALFF (15).

The associations between reported affective distress and ALFF values in limbic and striatal areas indicate that alterations in these areas may contribute to or reflect the ongoing emotional distress experienced by some trauma survivors. In fact, animal studies provide evidence that prefrontal-limbic and striatal systems play an important role in the regulation of stress responses. Such studies reveal that 30 to 60 min after the onset



**Fig. 2.** Decreased functional connectivity (blue line) mainly involving limbic-striatal areas and cerebellum in the survivor group compared with the control group ( $P < 0.05$ , corrected with FWE correction). ACC, anterior cingulate cortex; R-Insu, right insula; L-Insu, left insula; R-Cau, right caudate; L-Cau, left caudate; R-Amy, right amygdala; L-Amy, left amygdala; R-Hip, right hippocampus; L-Hip, left hippocampus; Cereb, cerebellum.



PNAS | September 8, 2009 | vol. 106 | no. 36 | 15415

dala, hippocampi, caudate, putamen, insula, precentral gyrus, and lateral parietal areas, and three medial areas (i.e., ACC, MPF, and PCC).

In SPM2, after bandpass filtering (0.01–0.08 Hz) (49) and linear-trend removal, a reference time series for each seed was extracted by averaging the fMRI time series of voxels within each ROI as defined in the WFU-pickatlas (Wake Forest University, Wake Forest, NC) (50, 51). Fifteen correlations were computed between each seed reference and the rest of the brain in a voxel-wise manner. Finally, the correlation coefficients in each voxel were transformed to  $z$  values by using the Fisher  $r$ -to- $z$  transformation to improve normality before averaging data across participants. By using SPM2, we removed components with high correlation to CSF or white matter, or with low correlation to gray matter, which are thought to be associated with artifacts, such as cardiac-induced (52) or respiratory-induced variations (53), rather than representing neural activity.

**Statistical Analysis.** ALFF maps in the survivor and control groups were compared on a voxelwise basis by using a two-sample  $t$  test in SPM2. First, a statistical threshold of  $P < 0.05$  (after FWE correction) was used for an exploratory whole-brain analysis. Second, SVC was performed to examine specific ROIs using a less conservative threshold ( $P < 0.05$  at cluster level). In ROIs, selected based on prior neuroimaging studies of PTSDs (5), we applied SVC by using masks (bilateral amygdala, hippocampi, caudate, putamen, and insula, as well as medial cingulate cortex) from the WFU-pickatlas (50, 51). SVC was also used in the four core areas of the default mode (19), including MPF, PCC, and bilateral lateral parietal areas.

ROI analysis was also used to explore the association between emotional

distress reflected by the SAS (20) and the SDS (21) with the average ALFF values in the areas of interest (Table 3). These were selected on the basis that they expressed ALFF alterations in the present investigation and/or were previously associated with altered emotional processing or episodic memory in patients with posttraumatic disorders (5). The average ALFF values of all voxels in these areas were extracted by using a volume of interest approach in SPM2 and were analyzed by using SPSS13.0. A multiple-correlation analysis was performed to estimate the relationship between the average ALFF values and SAS/SDS scores. Because of the lack of spatial dependency across the averaged time series from each ROI, a statistical threshold of  $P < 0.05$  (two-tailed) was used, with Bonferroni correction.

To examine functional connectivity within the groups, one-sample  $t$  tests were performed on the individual  $z$ -value maps with a statistical threshold of  $P < 0.05$  (after FWE correction). The  $z$  values in each voxel were compared across groups by using a two-sample  $t$  test in SPM2, again with statistical inferences being made at  $P < 0.05$  (after FWE correction). Finally, ROI analysis was also performed to characterize the association between functional connectivity and averaged regional ALFF, and between connectivity indices and reported level of emotional distress (SAS and SDS scores), by using a statistical threshold of  $P < 0.05$  with Bonferroni correction.

**ACKNOWLEDGMENTS.** We thank two anonymous reviewers for their multiple constructive comments. This study was supported by National Natural Science Foundation for Distinguished Young Scholars of China Grant 30625024, Programs from the State Education Ministry Grant SRFDP-20060610073, National Basic Research 973 Program 2007CB512305/2, and National High Technology Program of China 863 Program 2008AA022408.

- Perrin MA, et al. (2007) Differences in PTSD prevalence and associated risk factors among World Trade Center disaster rescue and recovery workers. *Am J Psychiatry* 164:1385–1394.
- Miller MM, McEwen BS (2006) Establishing an agenda for translational research on PTSD. *Ann NY Acad Sci* 1071:294–312.
- Jaferi A, Bhatnagar S (2006) Corticosterone can act at the posterior paraventricular thalamus to inhibit hypothalamic-pituitary-adrenal activity in animals that habituate to repeated stress. *Endocrinology* 147:4917–4930.
- Armory JL, Corbo V, Clement MH, Brunet A (2005) Amygdala response in patients with acute PTSD to masked and unmasked emotional facial expressions. *Am J Psychiatry* 162:1961–1963.
- Franconi V, Vermetten E, Bremner JD (2007) Functional neuroimaging studies in posttraumatic stress disorder: Review of current methods and findings. *Depress Anxiety* 24:202–218.
- Bryant RA, et al. (2008) Enhanced amygdala and medial prefrontal activation during nonconscious processing of fear in posttraumatic stress disorder: An fMRI study. *Hum Brain Mapp* 29:517–523.
- Stein MB, Simmons AN, Feinstein JS, Paulus MP (2007) Increased amygdala and insula activation during emotion processing in anxiety-prone subjects. *Am J Psychiatry* 164:318–327.
- Yang P, Wu MT, Hsu CC, Ker JH (2004) Evidence of early neurobiological alternations in adolescents with posttraumatic stress disorder: A functional MRI study. *Neurosci Lett* 370:13–18.
- Osuch EA, Willis MW, Bluhm R, Ursano RJ, Drevets WC (2008) Neurophysiological responses to traumatic reminders in the acute aftermath of serious motor vehicle collisions using [ $^{15}\text{O}$ ]-H $_2\text{O}$  positron emission tomography. *Biol Psychiatry* 64:327–335.
- Bryant RA, et al. (2005) Neural networks of information processing in posttraumatic stress disorder: A functional magnetic resonance imaging study. *Biol Psychiatry* 58:111–118.
- Driessen M, et al. (2004) Posttraumatic stress disorder and fMRI activation patterns of traumatic memory in patients with borderline personality disorder. *Biol Psychiatry* 55:603–611.
- Simmons AN, et al. (2008) Functional activation and neural networks in women with posttraumatic stress disorder related to intimate partner violence. *Biol Psychiatry* 64:681–690.
- Callicott JH, et al. (2003) Complexity of prefrontal cortical dysfunction in schizophrenia: More than up or down. *Am J Psychiatry* 160:2209–2215.
- Lui S, et al. (2008) Differential interictal activity of the precuneus/posterior cingulate cortex revealed by resting state functional MRI at 3T in generalized vs. partial seizure. *J Magn Reson Imaging* 27:1214–1220.
- Shmuel A, Leopold DA (2008) Neuronal correlates of spontaneous fluctuations in fMRI signals in monkey visual cortex: Implications for functional connectivity at rest. *Hum Brain Mapp* 29:751–761.
- Leopold DA, Murayama Y, Logothetis NK (2003) Very slow activity fluctuations in monkey visual cortex: Implications for functional brain imaging. *Cereb Cortex* 13:422–433.
- Laufs H, et al. (2003) Electroencephalographic signatures of attentional and cognitive default modes in spontaneous brain activity fluctuations at rest. *Proc Natl Acad Sci USA* 100:11053–11058.
- Goncalves SI, et al. (2006) Correlating the alpha rhythm to BOLD using simultaneous EEG/fMRI: Inter-subject variability. *NeuroImage* 30:203–213.
- Fox MD, et al. (2005) The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci USA* 102:9673–9678.
- Zung WW (1965) A self-rating depression scale. *Arch Gen Psychiatry* 12:63–70.
- Zung WW (1971) A rating instrument for anxiety disorders. *Psychosomatics* 12:371–379.
- Drevets WC (2000) Neuroimaging studies of mood disorders. *Biol Psychiatry* 48:813–829.
- Sakamoto H, et al. (2005) Parahippocampal activation evoked by masked traumatic images in posttraumatic stress disorder: A functional MRI study. *NeuroImage* 26:813–821.
- Cardinal RN, Parkinson JA, Hall J, Everitt BJ (2002) Emotion and motivation: The role of the amygdala, ventral striatum, and prefrontal cortex. *Neurosci Biobehav Rev* 26:321–352.
- Forstmann BU, et al. (2008) Striatum and pre-SMA facilitate decision-making under time pressure. *Proc Natl Acad Sci USA* 105:17538–17542.
- Lanius RA, et al. (2004) The nature of traumatic memories: A 4-T fMRI functional connectivity analysis. *Am J Psychiatry* 161:36–44.
- Ottowitz WE, et al. (2004) Neural and endocrine correlates of sadness in women: Implications for neural network regulation of HPA activity. *J Neuropsychiatry Clin Neurosci* 16:446–455.
- Damasio AR, et al. (2000) Subcortical and cortical brain activity during the feeling of self-generated emotions. *Nat Neurosci* 3:1049–1056.
- Morilak DA, et al. (2005) Role of brain norepinephrine in the behavioral response to stress. *Prog Neuropsychopharmacol Biol Psychiatry* 29:1214–1224.
- Pibiri F, Nelson M, Guidotti A, Costa E, Pinna G (2008) Decreased corticolimbic allopregnanolone expression during social isolation enhances contextual fear: A model relevant for posttraumatic stress disorder. *Proc Natl Acad Sci USA* 105:5567–5572.
- Lewis DA, Moghaddam B (2006) Cognitive dysfunction in schizophrenia: Convergence of gamma-aminobutyric acid and glutamate alterations. *Arch Neurol* 63:1372–1376.
- Lewis DA, et al. (2008) Subunit-selective modulation of GABA type A receptor neurotransmission and cognition in schizophrenia. *Am J Psychiatry* 165:1585–1593.
- Panksepp J, Nelson E, Bekkedal M (1997) Brain systems for the mediation of social separation-distress and social-reward. Evolutionary antecedents and neuropeptide intermediaries. *Ann NY Acad Sci* 807:78–100.
- Pacak K, et al. (1995) Decreased central and peripheral catecholaminergic activation in obese Zucker rats. *Endocrinology* 136:4360–4367.
- Etkin A, Wager TD (2007) Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 164:1476–1488.
- Strome EM, et al. (2002) Intracerebroventricular corticotropin-releasing factor increases limbic glucose metabolism and has social context-dependent behavioral effects in nonhuman primates. *Proc Natl Acad Sci USA* 99:15749–15754.
- Auer DP (2008) Spontaneous low-frequency blood oxygenation level-dependent fluctuations and functional connectivity analysis of the ‘resting’ brain. *Magn Reson Imaging* 26:1055–1064.
- Lanius RA, et al. (2005) Functional connectivity of dissociative responses in posttraumatic stress disorder: A functional magnetic resonance imaging investigation. *Biol Psychiatry* 57:873–884.
- Stevenson CW, Halliday DM, Marsden CA, Mason R (2007) Systemic administration of the benzodiazepine receptor partial inverse agonist FG-7142 disrupts corticolimbic network interactions. *Synapse* 61:646–663.
- Mason MF, et al. (2007) Wandering minds: The default network and stimulus-independent thought. *Science* 315:393–395.
- Greicius MD, Srivastava G, Reiss AL, Menon V (2004) Default-mode network activity distinguishes Alzheimer’s disease from healthy aging: Evidence from functional MRI. *Proc Natl Acad Sci USA* 101:4637–4642.

42. Garrity AG, et al. (2007) Aberrant "default mode" functional connectivity in schizophrenia. *Am J Psychiatry* 164:450–457.
43. Lanius RA, et al. (2009) Default mode network connectivity as a predictor of post-traumatic stress disorder symptom severity in acutely traumatized subjects. *Acta Psychiatr Scand*, in press.
44. Raichle ME, et al. (2001) A default mode of brain function. *Proc Natl Acad Sci USA* 98:676–682.
45. Horowitz SG, et al. (2008) Low frequency BOLD fluctuations during resting wakefulness and light sleep: A simultaneous EEG-fMRI study. *Hum Brain Mapp* 29:671–682.
46. Vincent JL, et al. (2007) Intrinsic functional architecture in the anaesthetized monkey brain. *Nature* 447:83–86.
47. Lui S, et al. (2009) Association of cerebral deficits with clinical symptoms in antipsychotic-naïve first-episode schizophrenia: An optimized voxel-based morphometry and resting state functional connectivity study. *Am J Psychiatry* 166:196–205.
48. Yang H, et al. (2007) Amplitude of low frequency fluctuation within visual areas revealed by resting-state functional MRI. *NeuroImage* 36:144–152.
49. Cordes D, et al. (2001) Frequencies contributing to functional connectivity in the cerebral cortex in "resting-state" data. *AJNR Am J Neuroradiol* 22:1326–1333.
50. Lancaster JL, et al. (2000) Automated Talairach atlas labels for functional brain mapping. *Hum Brain Mapp* 10:120–131.
51. Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH (2003) An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *NeuroImage* 19:1233–1239.
52. Dagli MS, Ingeholm JE, Haxby JV (1999) Localization of cardiac-induced signal change in fMRI. *NeuroImage* 9:407–415.
53. Windischberger C, et al. (2002) On the origin of respiratory artifacts in BOLD-EPI of the human brain. *Magn Reson Imaging* 20:575–582.